

The gaseous extent of galaxies and the origin of $\text{Ly}\alpha$ absorption systems. IV: $\text{Ly}\alpha$ absorbers arising in a galaxy group.¹

A. Ortiz-Gil^{2,3}, K.M. Lanzetta², J.K. Webb³, X. Barcons⁴, A. Fernández-Soto³

¹Based on observations with the NASA/ESA Hubble Space Telescope, obtained at the Space Telescope Science Institute, which is operated by the Association of Universities for Research in Astronomy, Inc., under NASA contract NAS5-26555.

²Department of Physics & Astronomy, State University of New York at Stony Brook, Stony Brook, NY 11794-3800, U.S.A.

³School of Physics, University of New South Wales, Kensington, NSW-2031, Sydney, AUSTRALIA

⁴Instituto de Física de Cantabria (Consejo Superior de Investigaciones Científicas–Universidad de Cantabria), 39005 Santander, SPAIN

ABSTRACT

We present new GHRS observations of Ly α absorption lines associated with a group of galaxies towards the QSO 1545+2101. We have identified eight distinct Ly α absorption features in the spectrum of QSO 1545+2101 at a mean redshift of $\langle z \rangle = 0.2648$ with a velocity dispersion of 163 km s $^{-1}$. A group of galaxies is detected in the vicinity of this QSO at a mean redshift of $\langle z \rangle = 0.2645$ and velocity dispersion 239 km s $^{-1}$.

The identification of discrete absorption systems indicates that they arise in clouds of neutral hydrogen rather than in a diffuse intragroup medium. Our analysis suggests that the Ly α absorption lines are associated with individual galaxies in the group, although a one-to-one relationship between absorbers and galaxies is difficult to establish in such a dense environment.

Subject headings: galaxies: clusters: general – galaxies: evolution – quasars: absorption lines

1. Introduction

$\text{Ly}\alpha$ absorption due to groups or clusters of galaxies has only been detected relatively recently (Lanzetta et al. 1996, Ortiz-Gil et al. 1997, Tripp et al. 1998, Shull et al. 1998). This is because it had been difficult to identify suitable combinations of background QSOs and foreground clusters close enough to the QSO line of sight to produce $\text{Ly}\alpha$ absorption in the QSO spectrum. The search for suitable groups was poorly motivated since $\text{Ly}\alpha$ absorption would not be expected due to the high temperature of the intracluster medium. Furthermore the majority of Lyman-alpha absorption line data in the literature are at redshifts $z > 1.5$ (when the line is shifted into the optical band), where most clusters are still on the process of formation.

Recent observations appear to produce contradictory results: Morris et al. (1993) and Bowen, Blades, & Pettini (1996) failed to identify absorption due to clusters of galaxies, in contrast to the results of Lanzetta, Webb & Barcons (1996), Ortiz-Gil et al. (1997), Tripp et al. (1998), and Shull et al. (1998).

However, other recent work suggests that galaxies may partially retain their gaseous envelopes in a cluster environment (see Cayatte et al. 1990 for an example in the Virgo cluster). Zabludoff & Mulchaey (1998) find that galaxies in poor groups ($\approx 20\text{--}50$ members) lie in a common halo which contains most of the mass of the group. Blitz et al. (1998) suggest that the High Velocity Clouds (HVCs) detected in the Local Group might be the counterparts of Lyman Limit systems, as they find similar column densities and internal velocity dispersions and subsolar metallicities. They also suggest that lower column density HVCs may correspond to $\text{Ly}\alpha$ clouds. In their models the HVCs trace the distribution of dark matter in and around the group, following its filamentary/sheet-like structure.

Lanzetta et al. (1996) report the identification of a group of galaxies toward QSO 1545+2101 responsible for a broad absorption feature present in an HST *Faint Object Spectrograph* (FOS) spectrum of this object. The spectral resolution of these data was insufficient to show whether it was the group as a whole or individual galaxies within it which were responsible for the observed $\text{Ly}\alpha$ absorption. Individual galaxies would give rise to discrete absorption components associated with particular galaxies. Also, galaxies at a smaller impact parameter should produce higher column densities.

In this paper we present new higher resolution observations of the QSO 1545+2101 using the *Goddard High Resolution Spectrograph* on HST.

Throughout this paper we have adopted a deceleration parameter $q_0 = 0.5$ and a dimensionless Hubble constant $h = H_0/100$ (km s⁻¹Mpc⁻¹).

2. HST/GHRS observations of the QSO 1545+2101

A high resolution spectrum of the QSO 1545+2101 was obtained using the GHRS spectrograph with the G160M grating on August 23 1996 (Fig. 1). The observations were obtained during a series of 14 exposures each of 300s duration for a total exposure time of 4200s. The individual exposures were reduced using standard pipeline techniques and were registered to a common origin and combined using our own reduction programs. The final spectrum was fitted with a smooth continuum using an iterative spline fitting technique. The spectral resolution of the final spectrum was measured to be $\text{FWHM}=0.07 \text{ \AA}$ (or $\text{FWHM}=14 \text{ km s}^{-1}$), and the continuum signal-to-noise ratio was measured to be $S/N \approx 12$ per resolution element. Previous imaging and spectroscopic observations of the field surrounding QSO 1545+2101 are described in Lanzetta et al. (1996).

3. Detection and analysis of the absorption systems in the field of QSO 1545+2101

We detected all absorption lines above a significance level of 3σ in the spectrum of QSO 1545+2101. The parameters characterizing each absorption line were estimated using multi-component Voigt profile fitting (three different and independently developed software packages were used, the three of them providing the same results). The values obtained are shown in Table 1. The 3σ confidence level detection limit was found to be $0.03 \text{ \AA}$.

Two absorption lines are found at $z = 0.2504707 \pm 0.0000030$ and $z = 0.2522505 \pm 0.0000016$, with a velocity separation of $\sim 427 \text{ km s}^{-1}$. A group of at least eight lines is also detected, the redshift centroid of this group being $\langle z \rangle = 0.2648 \pm 0.0002$ with a velocity dispersion of $163 \pm 57 \text{ km s}^{-1}$.

Galactic heavy element lines in the spectrum (C IV $\lambda\lambda 1548, 1550$ and Si II $\lambda 1526$) were identified and removed them from subsequent analyses (Fig.1).

4. Group of galaxies in the field of QSO 1545+2101

The spectroscopic galaxy sample was selected on the basis of galaxy brightness and proximity to the QSO line-of-sight. Although this sample cannot be considered complete the galaxies were selected randomly from an essentially flux-limited sample. The sample of galaxies used in this study is given in Table 2. Figure 2 is an image of the field. The galaxies in the group which were observed spectroscopically are indicated in that figure. The error

in the galaxies redshift is estimated to be $\sim 120 \text{ km s}^{-1}$ (Lanzetta et al. 1995).

We detected a group of seven galaxies with redshift centroid at $\langle z \rangle = 0.2645 \pm 0.0004$ and velocity dispersion of $239 \pm 90 \text{ km s}^{-1}$. Their individual impact parameters range from 7.2 up to $456.4 h^{-1} \text{ kpc}$.

We searched in the archives for X-ray data on this field as detecting X-ray emission from the group might help to characterize it better. Unfortunately the group is so close to the QSO (which is an X-ray source itself) that only the *ROSAT* High Resolution Imager data would be of any use, and there are no such data in the archive. QSO 1545+2101 was observed both with the *Einstein* Imaging Proportional Counter and with the *ROSAT* Position Sensitive Proportional Counter, but the extended Point-Spread-Function of both instruments resulted in QSO emission severely contaminating the region where the galaxy group might emit X-rays.

In any case, from our image of this field and the fact that the velocity dispersion that we have measured for this group is quite small, it is clear that it is a loose association of galaxies rather than a galaxy cluster. The maximum value of impact parameter in our sample ($456.4 h^{-1} \text{ kpc}$) is also the typical physical size for poor groups of galaxies (Zabludoff & Mulchaey 1998). In fact, this group is most probably the one hosting the QSO itself, as $z_{em} = 0.264 \pm 0.0003$ for this object (Marziani et al. 1996). One might therefore expect this group to have only between 10-20 members (Bahcall et al. 1997).

5. The relationship between the absorption systems and the galaxies

The cluster of absorption lines present in the spectrum of QSO 1545+2101 might arise in different environments. They might be intrinsic absorbers, arising either in the QSO region itself or in its near environment. The similar redshifts of the QSO and the group of absorbers in these data may point to one of these hypotheses as the right one. In addition, QSO 1545+2101 is a radio-loud object and there have been suggestions that intrinsic absorption or absorption arising in the QSO host galaxy would be stronger in radio-loud QSOs than in radio-quiet ones (Foltz et al. 1988; Mathur, Wilkes & Aldcroft 1997).

But there are also some arguments against an intrinsic origin for these lines, as discussed by Lanzetta et al. (1996). Associated absorbers of the kind described above produce rather strong absorption lines, while the lines detected in QSO 1545+2101 are relatively weak. Moreover, no corresponding metal absorption lines have been detected. With the new data from GHRS we have more evidence against the associated nature of the absorption systems: the agreement between the galaxies' and absorbers' redshift centroids and between their

corresponding velocity dispersions suggests that both galaxies and absorbers groups share—at least—the same physical location. These characteristics lead us to reject the associated hypothesis and to consider another possible scenario: absorption arising in cosmologically intervening objects, within the same group of galaxies that hosts the QSO.

To further explore this third hypothesis a demonstration of the non-random coincidence between the galaxies and absorbers positions in velocity space is necessary. Identifying each galaxy with a single absorption line would also be very interesting. In what follows we use statistical methods to address both questions.

5.1. A cluster of absorbers arising in a group of galaxies

The group of galaxies detected toward the QSO 1545+2101 has a mean redshift of $\langle z_g \rangle = 0.2645 \pm 0.0004$, compatible with the mean redshift centroid value of the group of absorption lines, $\langle z_a \rangle = 0.2648 \pm 0.0002$ (to the red of the Galactic Si II line in Fig. 1). The absorber and galaxy velocity dispersions are similar: $239 \pm 90 \text{ km s}^{-1}$ for the group of galaxies and $163 \pm 57 \text{ km s}^{-1}$ for the group of absorption lines (the error in the galaxy redshifts is $\sim 120 \text{ km s}^{-1}$). This strongly suggests a connection between the absorbers and galaxies.

There is also a galaxy in this field whose redshift is $z = 0.2510$. Two Ly α absorption lines are detected near this redshift at $z = 0.2504707 \pm 0.0000030$ and $z = 0.2522505 \pm 0.0000020$ (to the blue of the Galactic Si II line in Fig. 1). The galaxy-absorber velocity differences are $\Delta v = 127 \pm 120 \text{ km s}^{-1}$ and $\Delta v = 300 \pm 120 \text{ km s}^{-1}$ respectively. This implies that this galaxy, whose impact parameter is $\rho = 306.4 \text{ h}^{-1} \text{ kpc}$, could be responsible for one of the absorption lines as both Δv values are compatible with the velocity dispersions that one finds typically in a galactic halo ($\approx 200 \text{ km s}^{-1}$). The velocity difference between the two absorption systems (427 km s^{-1}), is perhaps too large for the same galaxy to be responsible for both of them. It may also be that we have not observed the actual galaxy giving rise to either of these absorption lines, since our galaxy sample is not complete.

5.1.1. Statistical analysis

Two statistical tests were carried out to investigate the relationship between the group of galaxies and absorbers. In the first, we computed the two-point cross-correlation function (ξ_{ag}) between the absorbers and the galaxies. This function was normalized by computing the ξ_{ag} expected if there were no relation between absorbers and galaxies (derived using

galaxy redshifts which are randomly distributed over a redshift range around the real group of absorption lines). Errors were computed using a bootstrap method, simulating 1000 samples of 7 galaxy redshifts, each set randomly selected from the real set, deriving error estimates from the distribution of 1000 values of ξ_{ag} . The final result is shown in Fig. 3 (panel *a*).

In the second statistical test, we applied a similar statistical method, not to the actual set of absorbers but to the individual pixel intensities in the spectrum. This way we overcome any potential problems introduced as a consequence of incorrect determination of the true velocity structure in the profile fitting process, or the presence of weak lines falling below the detection threshold. Recent analyses by Liske, Webb & Carswell (1999) show that the study of pixel intensities is more sensitive to clustering than the usual line-fitting techniques. The test is as follows: we evaluated for each pixel i the value of the function $g_i = 1 - f_i$, where f_i is the intensity of pixel i normalized to the continuum. Clearly, g_i has larger values in pixels belonging to absorption lines. For pixels corresponding to metal lines (including galactic lines) we assigned a value of $g_i = 0$. Then, for each galaxy we compute the function $\zeta_v = \sum_{n=1}^N \sum_j g_{nj}$, where g_{nj} refers to all the pixels j located at a distance v in velocity space from galaxy n , N being the total number of galaxies in the sample. The velocity distances v considered range from 0 km s⁻¹ up to the one spanned by the whole spectrum. This function ζ_v is analogous in some sense to the two-point cross-correlation function computed before: high values of ζ_v at low velocity distances reflect the tendency of the pixels corresponding to absorption lines to lie close to the galaxies. The same function ζ_v corresponding, again, to a randomly selected sample of galaxies chosen from a uniformly distributed group was computed in a way analogous to the previous one and the result, after normalizing to this random case, is shown in Fig. 3 (panel *b*). The error bars in ζ_v were computed using a bootstrap method as before. The result obtained is similar to the one obtained by computing the two-point cross-correlation function: most of the pixels belonging to absorption lines (i.e., with larger values of g) lie less than 200 km s⁻¹ away from the galaxies.

5.2. Are absorption lines related to galaxies on a case-by-case basis?

As there is a clear complex of discrete Ly α lines in the spectrum of QSO 1545+2101, we explored the possibility of a one-to-one match between galaxies and absorbers.

A Gaussian model was assumed for the distribution of galaxies in the group. The null hypothesis is that the galaxies are randomly drawn from a Gaussian whose parameters are derived from the real data. The two-point correlation functions ξ_{ag} and the function ζ_v were computed in the same way as in §5.1 (see Fig. 3, panels *c* and *d*). No evidence of a one-to-one

correspondence between galaxies and absorbers is found.

We can estimate the maximum statistical velocity dispersion between individual absorbers and their galaxies of origin that would permit the detection of a one-to-one correspondence, assuming an intrinsic one-to-one correspondence exists. A Kolmogorov-Smirnov Monte Carlo test showed that if the average velocity dispersion between the galaxies and the corresponding absorbers were $\lesssim 100 \text{ km s}^{-1}$ then the null hypothesis would be rejected ($> 2\sigma$) in 90% of the cases. As a typical galactic velocity dispersion is $\sim 200 \text{ km s}^{-1}$, this condition is not likely to be satisfied in practice. Another possibility is to improve the statistics. From Monte Carlo simulations we estimate that about 100 groups similar to the ones studied here are needed for a 2σ detection of this one-to-one association, for a velocity dispersion between the absorber and the galaxy of about 200 km s^{-1} .

6. Discussion and conclusions

We have detected a clump of absorption lines along the line-of-sight towards the QSO 1545+2101. A group of galaxies has also been detected, with impact parameters ρ of the individual galaxies to the QSO line-of-sight of less than $\sim 460h^{-1} \text{ kpc}$. The group is probably the one hosting the QSO, so we can expect it to have about 10-20 members (Bahcall et al. 1997).

Several scenarios might give rise to the absorption. Due to the close redshift values of the QSO and the group of absorbers one could easily think that they arise either in the QSO itself or in the corresponding host galaxy. We consider that there are compelling arguments supporting the intervening system hypothesis (see Lanzetta et al. 1996) and contradicting the associated hypothesis. We now summarise those arguments.

The velocity spanned by the group of $\text{Ly}\alpha$ absorption lines is consistent with the velocity dispersion of the group of galaxies. This implies that the $\text{Ly}\alpha$ absorbers arising in that group occupy the same region of space as the galaxies themselves. Moreover, the spectrum of QSO 1545+2101 reveals a group of *discrete* $\text{Ly}\alpha$ absorption lines at $z \approx 0.26$. Multi-component Voigt profile fitting provides a statistically good fit to the data, indicating that the absorption lines arise in overdense gas regions rather than in some smoothly distributed intragroup medium.

The average Doppler dispersion parameter of the absorption lines, b , is measured to be $19 \pm 4 \text{ km s}^{-1}$ with a dispersion of $10 \pm 4 \text{ km s}^{-1}$. This value is in agreement with the values measured in the low redshift $\text{Ly}\alpha$ forest. Therefore there is no evidence from this case for any physical difference, in terms of the b parameter, between $\text{Ly}\alpha$ clouds lying within or

outside of groups.

Two statistical analyses show that the distribution of galaxies with respect to the absorbers is not random, but that is not possible to confirm a one-to-one match due to the proximity in velocity space of the galaxies in the groups and the uncertainty on their redshifts. A Kolmogorov-Smirnov test showed that a small galaxy-absorber velocity dispersion (less than $\sim 100 \text{ km s}^{-1}$) would be required to establish a one-to-one match. As this is well below the typical values corresponding to a galaxy potential well, another approach is required, such as having a large enough sample of clusters or groups of galaxies related to clusters of Ly α absorption lines. All the facts above support the idea of a physical connection between the group of galaxies and the group of Ly α absorbers.

Another piece of circumstantial evidence pointing to a one-to-one relationship between absorbers and galaxies. This concerns the number of each type of object detected. As mentioned before, the galaxy group towards QSO 1545+2101 contains approximately 10-20 members. According to Lanzetta et al. (1995), only a subset of them will be close enough to the QSO line of sight to produce observable Ly α absorption ($\rho \lesssim 160 \text{ kpc}$ for a covering factor of ≈ 1). We observe eight individual components in the absorption profile, consistent with the expectations from such a naïve model. There is no obvious reason why such an agreement would be found for some other quite different model (be it HVCs, filaments or any other structure). We note that the absorption lines may break up into further components at higher spectral resolution although these may then be substructure within individual galaxies.

A.O.-G., K.M.L. and A.F.-S. were supported by grant NAG-53261; grants AR-0580-30194A, GO-0594-80194, GO-0594-90194A and GO-0661-20195A from STScI; and grant AST-9624216 from NSF. A.O.-G. acknowledges support from a UNSW Honorary Fellowship. A.F.-S. was also supported by an ARC grant. X.B. was partially supported by the DGES under project PB95-0122.

REFERENCES

- Bahcall, J.N., Kirhakos, S., Saxe, D.H., & Schneider, D.P. 1997, *ApJ*, 479, 642
- Blitz, L., Spergel, D.N., Teuben, P.J., Hartmann, D., & Burton, W.B. 1998, *ApJ* submitted (also [astro-ph/9803251](#))
- Bowen, D. V., Blades, J. C., & Pettini, M. 1996, *ApJ*, 464, 141
- Cayatte, V., Van Gorkom, J. H., Balkowski, C., & Kotanyi, K. 1990, *AJ*, 100, 604
- Foltz, C. B., Chaffe, F. H., Weymann, R. J., & Anderson, S. F. 1988 in *QSO Absorption Lines*, ed J. C. Blades, D. Turnshek, & C. Norman (Cambridge University Press), 53
- Lanzetta, K. M., Bowen, D. V., Tytler, D., & Webb, J. K. 1995, *ApJ*, 442, 538
- Lanzetta, K. M., Webb, J. K., & Barcons, X. 1996, *ApJ*, 456, L17
- Liske, J., Webb, J. K., & Carswell, R. F. 1999, *MNRAS*, 301, 787
- Marziani, P., Sulentic, J. W., Dultzin-Hacyan, D., Calvani, M., & Moles, M. 1996, *ApJS*, 104, 37
- Mathur, S., Wilkes, B.J., Aldcroft, T. 1997, *ApJ*, 478, 182
- Morris, S. L., Weymann, R. J., Dressler, A., McCarthy, P. J., Smith, B. A., Terrile, R. J., Giovanelli, R., & Irwin, M. J. 1993, *ApJ*, 419, 524
- Ortiz-Gil, A., Lanzetta, K. M., Webb, J. K., & Barcons, X. 1997, *Proceedings of the XIII IAP Colloquium (Paris)*, 438
- Shull, J. M., Penton, S. V., Stocke, T. J., Giroux, M. L., van Gorkom, J. H., Lee, Y. -H., & Carilli, C. 1998, *AJ*, 116, 2094
- Tripp, T. M., Lu, L., & Savage, B. 1998, *ApJ*, 508, 200
- Zabludoff, A. I., & Mulchaey, J. S. 1998, *ApJ*, 496, 39

Fig. 1.— Spectrum of QSO 1545+2101, smoothed at the Nyquist rate. The original data have a resolution of $\text{FWHM} = 0.07\text{\AA}$ ($\text{FWHM} = 14 \text{ km s}^{-1}$). Tick marks in the upper panel indicate the predicted wavelengths of $\text{Ly}\alpha$ at the redshifts of the galaxies listed in Table 2. Tick marks in the lower panel show the positions of the detected absorption lines

Fig. 2.— Image of a $\approx 12' \times 12'$ field toward QSO 1545+2101 (north is up, east is left). Marked galaxies are the ones belonging to the group. Numbers increase with increasing z . The QSO is marked by number 0

Fig. 3.— (a): cross-correlation function between galaxies and absorbers normalized using a uniform distribution. The existence of a correlation between them is clear. (b): normalized ζ_v function (see text). There is a clear connection between the real sample of galaxies and the pixels corresponding to absorption lines. (c): cross-correlation function between galaxies and absorbers normalized using a Gaussian distribution. No significant correlation between them is found. (d): normalized ζ_v function (see text). A one-to-one match between the absorption systems and the galaxies cannot be established

z_{abs}	σ_z (10^{-7})	W ($\text{\AA}$)	$\sigma(W)$ ($10^{-4} \text{\AA}$)	b (km s^{-1})	$\sigma(b)$ (km s^{-1})	$\log N$ (cm^{-2})	$\sigma(\log N)$ (cm^{-2})
0.2504707	30	0.067	0.013	66	9	13.61	0.05
0.2522505	16	0.21	0.03	40	5	13.63	0.04
0.2634336	3	0.387	0.018	28.8	1.4	14.17	0.04
0.2641748	3	0.307	0.016	28.0	1.2	13.94	0.02
0.2645921	4	0.294	0.015	27.3	1.3	13.91	0.02
0.2648250	6	0.033	0.009	4	5	12.84	0.12
0.2650173	10	0.254	0.015	26	3	13.81	0.05
0.2652287	23	0.117	0.014	23	6	13.33	0.12
0.2654554	9	0.041	0.012	11	4	12.84	0.08
0.2656921	4	0.203	0.017	23.4	1.5	13.67	0.03

Table 1: Ly α absorption systems toward QSO 1545+2101. For each absorption line, z_{abs} is the redshift, W is the rest-frame equivalent width, b is the Doppler parameter and N is the column density

$\Delta\alpha$ (arcsec)	$\Delta\delta$ (arcsec)	θ (arcsec)	R	z_{gal}	ρ (h^{-1} kpc)	$M_R - 5 \log h$
-38.7	-118.3	124.5	20.1	0.2510	306.4	-19.4
58.3	31.2	66.1	19.2	0.2630	167.7	-20.4
53.4	-171.5	179.6	20.2	0.2638	456.4	-19.4
7.0	166.5	166.6	20.6	0.2639	423.5	-19.0
16.6	-8.7	18.7	19.9	0.2639	47.6	-19.7
85.2	-81.3	117.8	18.8	0.2652	300.2	-20.8
-2.7	-1.1	2.9	19.1	0.2657	7.2	-20.5
-21.8	98.8	101.2	18.8	0.2658	258.3	-20.8

Table 2: Group of galaxies in the field of QSO 1545+2101. For each galaxy, $\Delta\alpha$ and $\Delta\delta$ are the offset from the QSO in right ascension and declination, θ is the angular offset from the QSO, R is the observed R -band magnitude, z_{gal} is the redshift, ρ is the impact parameter to the QSO line of sight and M_R is the absolute R -band magnitude




Fig. 1.—




Fig. 2.—




Fig. 3.—